

FALL 2003 DATA FOR COCKTOWN CREEK

| Species | Site 1 to 2 | Site 2 to 3 | Site 3 to 4 | Site 4 to 5 | Site 5 to 6 | Site 6 to 7 | Site 7 to 8 | Site 8 to 9 | Site 9 to 10 | Total SAV observed | Frequency (In 52 Surveys) | % of total SAV found |
|------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|--------------|--------------------|---------------------------|----------------------|
| Coontail | 5 | 6 | 6 | 6 | 6 | 5 | 6 | 6 | 6 | 52 | 100% | 20% |
| Curly Pondweed | 4 | 3 | 4 | 6 | 5 | 6 | 6 | 6 | 6 | 46 | 88% | 17% |
| Wild Celery | 4 | 1 | 2 | 1 | 2 | 4 | 6 | 6 | 6 | 32 | 62% | 12% |
| Common Waterweed | 6 | 5 | 5 | 6 | 5 | 5 | 5 | 5 | 6 | 48 | 92% | 18% |
| Slender Pondweed | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 1 | 3 | 6 | 12% | 2% |
| Hydrilla | 6 | 6 | 5 | 6 | 6 | 5 | 6 | 6 | 4 | 51 | 98% | 19% |
| Horned Pondweed | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0% |
| Naiad | 3 | 3 | 4 | 4 | 3 | 3 | 3 | 2 | 3 | 28 | 54% | 11% |
| Southern Naiad | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0% |
| Species "richness" | 6 | 6 | 7 | 6 | 7 | 6 | 6 | 7 | 7 | | | |
| Total Observed at Site | 28 | 24 | 27 | 29 | 28 | 29 | 32 | 32 | 34 | 263 | | |
| No SAV found | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0% | |