

FALL 2006 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 36 Surveys)	% of total SAV found
Coontail	1	4	3	4	3	1	3	4	4	27	75%	29%
Curly Pondweed	0	0	0	0	1	0	1	2	3	7	19%	7%
Wild Celery	0	0	0	0	0	0	2	1	2	5	14%	5%
Common Waterweed	0	0	2	0	0	0	0	0	0	2	6%	2%
Slender Pondweed	0	0	1	0	0	0	0	0	0	1	3%	1%
Hydrilla	4	4	4	4	4	4	4	4	4	36	100%	38%
Horned Pondweed	0	0	0	0	0	0	0	0	0	0	0%	0%
Naiad	0	3	3	3	2	3	1	0	1	16	44%	17%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	2	3	5	3	4	3	5	4	5			
Total Observed at Site	5	11	13	11	10	8	11	11	14	94		
No SAV found	0	0	0	0	0	0	0	0	0		0%	