

## FALL 2008 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 65 Surveys)	% of total SAV found
Coontail	2	1	3	2	2	3	4	6	6	29	45%	18%
Curly Pondweed	1	0	0	0	0	1	1	2	6	11	17%	7%
Wild Celery	0	0	0	0	0	1	1	7	5	14	22%	9%
Common Waterweed	5	0	0	0	0	1	0	0	1	7	11%	4%
Slender Pondweed	0	0	0	0	0	0	0	0	0	0	0%	0%
Hydrilla	7	8	7	7	7	8	8	7	6	65	100%	41%
Horned Pondweed	0	0	0	0	0	0	0	0	0	0	0%	0%
Naiad	4	6	4	4	5	4	2	1	3	33	51%	21%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	5	3	3	3	3	6	5	5	6			
Total Observed at Site	24	15	14	13	14	24	16	23	23	159		
No SAV found	0	0	0	0	0	0	0	0	0		0%	