

## FALL 2009 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 57 Surveys)	% of total SAV found
Coontail	2	6	4	4	1	4	3	4	4	32	56%	30%
Curly Pondweed	0	0	0	0	0	0	1	0	3	4	7%	4%
Wild Celery	0	0	0	0	0	1	1	5	3	10	18%	9%
Common Waterweed	0	0	0	1	0	0	0	0	2	3	5%	3%
Slender Pondweed	0	0	0	0	0	0	1	0	0	1	2%	1%
Hydrilla	11	8	7	7	5	7	3	5	4	57	100%	53%
Horned Pondweed	0	0	0	0	0	0	0	0	0	0	0%	0%
Naiad	1	4	2	1	1	1	0	0	0	10	18%	9%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	3	3	3	4	3	4	5	3	5			
Total Observed at Site	14	18	13	13	7	13	9	14	16	107		
No SAV found	0	0	0	0	0	0	0	0	0		0%	