

SPRING 1999 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 26 Surveys)	% of total SAV found
Coontail	1	1	1	0	2	0	2	2	2	11	42%	13%
Curly Pondweed	2	2	3	2	3	2	3	3	3	23	88%	27%
Wild Celery	0	1	0	1	2	1	3	3	1	12	46%	14%
Common Waterweed	0	0	1	1	1	1	1	1	3	9	35%	11%
Slender Pondweed	1	2	2	2	0	1	1	2	1	12	46%	14%
Hydrilla	0	0	0	1	0	0	0	0	0	1	4%	1%
Horned Pondweed	1	2	3	1	3	1	1	2	2	16	62%	19%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	4	5	5	5	5	4	6	6	6			
Total Observed at Site	5	8	10	8	11	6	11	13	12	84		
No SAV found	0	0	0	0	0	0	0	0	0		0%	