

FALL 2000 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 62 Surveys)	% of total SAV found
Coontail	4	4	4	4	5	5	6	6	7	45	73%	16%
Curly Pondweed	3	2	5	4	3	5	6	7	7	42	68%	15%
Wild Celery	4	4	4	6	6	7	7	7	7	52	84%	19%
Common Waterweed	7	7	7	7	7	7	7	6	7	62	100%	23%
Slender Pondweed	1	1	1	2	0	2	0	1	0	8	13%	3%
Hydrilla	1	0	3	0	1	3	2	0	0	10	16%	4%
Horned Pondweed	2	0	1	0	0	0	0	0	0	3	5%	1%
Naiad	7	7	7	7	6	6	5	7	1	53	85%	19%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	8	6	8	6	6	7	6	6	5			
Total Observed at Site	29	25	32	30	28	35	33	34	29	275		
No SAV found	0	0	0	0	0	0	0	0	0		0%	