

SPRING 2002 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 54 Surveys)	% of total SAV found
Coontail	1	4	3	4	5	6	6	6	6	41	76%	23%
Curly Pondweed	3	4	4	5	4	6	4	6	5	41	76%	23%
Wild Celery	1	1	2	2	2	3	3	2	2	18	33%	10%
Common Waterweed	0	2	1	2	1	5	1	3	4	19	35%	10%
Slender Pondweed	1	0	0	0	1	0	0	0	0	2	4%	1%
Hydrilla	1	1	1	1	0	2	3	1	1	11	20%	6%
Horned Pondweed	4	6	5	5	5	6	5	6	6	48	89%	27%
Naiad	0	0	1	0	0	0	0	0	0	1	2%	1%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	6	6	7	6	6	6	6	6	6			
Total Observed at Site	11	18	17	19	18	28	22	24	24	181		
No SAV found	0	0	0	0	0	0	0	0	0		0%	