

SPRING 2003 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 64 Surveys)	% of total SAV found
Coontail	3	5	5	7	5	6	6	6	7	50	78%	19%
Curly Pondweed	4	2	6	8	6	6	6	7	7	52	81%	20%
Wild Celery	0	1	1	2	1	4	2	3	5	19	30%	7%
Common Waterweed	2	4	2	6	4	5	4	5	5	37	58%	14%
Slender Pondweed	4	2	4	5	3	4	4	5	6	37	58%	14%
Hydrilla	1	0	0	2	1	2	2	2	2	12	19%	5%
Horned Pondweed	7	7	6	8	6	6	5	5	5	55	86%	21%
Naiad	0	0	0	0	1	0	0	0	0	1	2%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	6	6	6	7	8	7	7	7	7			
Total Observed at Site	21	21	24	38	27	33	29	33	37	263		
No SAV found	0	0	0	0	0	0	0	0	0		0%	