

SPRING 2004 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 51 Surveys)	% of total SAV found
Coontail	3	5	3	8	6	7	6	6	6	50	88%	17%
Curly Pondweed	5	6	6	8	6	7	6	6	6	56	98%	19%
Wild Celery	0	1	1	1	3	3	2	1	2	14	25%	5%
Common Waterweed	4	4	3	7	4	6	6	6	6	46	81%	15%
Slender Pondweed	3	4	4	7	6	7	6	5	6	48	84%	16%
Hydrilla	2	3	1	5	3	5	3	1	2	25	44%	8%
Horned Pondweed	6	6	6	8	6	7	6	6	6	57	100%	19%
Naiad	0	0	1	1	0	0	0	0	1	3	5%	1%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	6	7	8	8	7	7	7	6	8			
Total Observed at Site	23	29	25	45	34	42	35	31	35	299		
No SAV found	0	0	0	0	0	0	0	0	0		0%	