

SPRING 2005 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 77 Surveys)	% of total SAV found
Coontail	3	3	4	4	4	3	3	3	4	21	58%	15%
Curly Pondweed	2	3	4	3	3	4	3	4	4	30	83%	21%
Wild Celery	0	0	0	1	0	0	1	0	0	2	6%	1%
Common Waterweed	3	2	3	1	2	3	3	3	4	24	67%	17%
Slender Pondweed	4	2	3	3	3	3	4	3	2	27	75%	19%
Hydrilla	2	0	0	0	0	2	1	1	1	7	19%	5%
Horned Pondweed	4	4	3	4	4	3	4	4	2	32	89%	22%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	6	5	5	6	5	6	7	6	6			
Total Observed at Site	18	14	17	16	16	18	19	24	17	143		
No SAV found	0	0	0	0	0	0	0	0	0		0%	