

SPRING 2006 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 19 Surveys)	% of total SAV found
Coontail	1	2	2	2	2	2	1	2	2	16	84%	20%
Curly Pondweed	0	2	2	2	2	2	1	2	2	15	79%	19%
Wild Celery	1	2	1	0	0	1	0	1	0	6	32%	7%
Common Waterweed	0	0	0	0	1	0	0	0	1	2	11%	2%
Slender Pondweed	1	1	2	1	1	0	0	1	0	7	37%	9%
Hydrilla	1	3	2	2	2	1	1	2	2	16	84%	20%
Horned Pondweed	2	3	2	2	2	2	2	2	2	19	100%	23%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	5	6	6	5	6	3	4	6	5			
Total Observed at Site	6	13	11	9	10	8	5	10	14	81		
No SAV found	0	0	0	0	0	0	0	0	0		0%	