

SPRING 2007 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 50 Surveys)	% of total SAV found
Coontail	1	1	3	4	3	2	1	5	4	24	48%	17%
Curly Pondweed	0	0	2	3	5	3	3	4	5	25	50%	18%
Wild Celery	0	1	1	0	0	2	1	1	1	7	14%	5%
Common Waterweed	0	0	0	2	2	0	2	3	1	10	10%	20%
Slender Pondweed	0	0	0	1	0	0	1	0	1	3	6%	2%
Hydrilla	3	1	4	3	3	2	5	3	4	28	56%	20%
Horned Pondweed	5	5	5	5	5	5	4	5	3	42	84%	30%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	2	3	5	3	4	3	5	4	5			
Total Observed at Site	9	8	15	18	18	14	17	21	19	139		
No SAV found	0	0	0	0	0	0	0	0	0		0%	