

## SPRING 2008 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 49 Surveys)	% of total SAV found
Coontail	0	1	3	5	3	3	4	5	5	29	59%	21%
Curly Pondweed	1	0	4	1	3	3	4	5	5	26	53%	19%
Wild Celery	0	1	1	0	0	0	0	0	0	2	4%	1%
Common Waterweed	0	0	1	0	0	0	0	0	2	3	6%	2%
Slender Pondweed	1	1	1	0	0	0	1	0	2	3	6%	2%
Hydrilla	1	2	4	3	3	2	3	4	4	26	53%	19%
Horned Pondweed	6	6	6	5	5	5	4	5	5	47	96%	34%
Naiad	0	0	0	0	0	0	1	0	0	1	2%	1%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	4	5	7	4	4	4	6	4	6			
Total Observed at Site	13	16	27	14	14	13	23	19	29	140		
No SAV found	0	0	0	0	0	0	0	0	0		0%	