

SPRING 2009 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 35 Surveys)	% of total SAV found
Coontail	1	3	2	2	1	1	1	3	3	17	49%	17%
Curly Pondweed	1	2	2	1	2	1	0	3	3	15	43%	15%
Wild Celery	0	0	0	1	0	0	0	1	1	3	9%	3%
Common Waterweed	0	0	0	0	1	0	0	0	1	2	6%	2%
Slender Pondweed	1	1	0	0	0	0	1	2	2	7	20%	7%
Hydrilla	3	3	2	4	2	2	2	2	1	21	60%	21%
Horned Pondweed	5	5	4	5	3	4	3	3	3	35	100%	35%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	5	5	4	5	5	4	4	6	7			
Total Observed at Site	11	14	10	13	9	8	7	14	14	100		
No SAV found	0	0	0	0	0	0	0	0	0		0%	