

SPRING 2010 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 44 Surveys)	% of total SAV found
Coontail	1	0	5	4	3	5	4	4	2	28	64%	18%
Curly Pondweed	0	1	3	2	1	4	4	3	3	21	48%	14%
Wild Celery	0	0	1	1	0	1	0	0	0	3	7%	2%
Common Waterweed	0	0	1	1	0	0	0	1	2	5	11%	3%
Slender Pondweed	0	0	3	4	1	1	3	2	2	16	36%	10%
Hydrilla	5	5	5	6	4	5	4	5	2	41	93%	26%
Horned Pondweed	5	4	5	5	4	6	4	5	3	41	93%	26%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	3	2	5	6	3	5	5	6	6			
Total Observed at Site	11	10	23	23	13	22	19	20	14	155		
No SAV found	0	0	0	0	0	0	0	0	0		0%	