

SPRING 2012 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 48 Surveys)	% of total SAV found
Coontail	2	3	3	4	5	6	4	6	5	38	79%	25%
Curly Pondweed	0	0	0	0	2	2	0	1	5	10	21%	7%
Wild Celery	0	0	0	0	0	0	0	2	1	3	6%	2%
Common Waterweed	0	0	0	0	0	0	0	0	0	0	0%	0%
Slender Pondweed	1	0	2	1	0	3	2	1	3	13	27%	9%
Hydrilla	3	6	6	5	6	5	6	6	5	47	98%	31%
Horned Pondweed	4	6	5	2	4	6	4	6	4	41	85%	29%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	4	3	4	4	4	5	4	6	6			
Total Observed at Site	10	15	16	12	16	23	15	22	23	152		
No SAV found	0	0	0	0	0	0	0	0	0		0%	