

SPRING 2013 DATA FOR COCKTOWN CREEK

Species	Site 1 to 2	Site 2 to 3	Site 3 to 4	Site 4 to 5	Site 5 to 6	Site 6 to 7	Site 7 to 8	Site 8 to 9	Site 9 to 10	Total SAV observed	Frequency (In 45 Surveys)	% of total SAV found
Coontail	0	0	2	2	2	2	3	5	5	21	48%	22%
Curly Pondweed	0	0	0	0	0	0	1	1	2	4	9%	4%
Wild Celery	0	0	0	1	0	0	0	0	1	2	5%	2%
Common Waterweed	0	0	0	0	0	0	0	0	0	0	0%	0%
Slender Pondweed	0	0	0	1	0	1	0	2	4	8	18%	8%
Hydrilla	0	1	4	3	2	1	2	4	3	20	45%	21%
Horned Pondweed	3	5	6	4	3	4	5	6	5	41	93%	43%
Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Southern Naiad	0	0	0	0	0	0	0	0	0	0	0%	0%
Species "richness"	1	2	3	4	3	3	4	5	6			
Total Observed at Site	3	6	12	11	7	8	11	18	20	96		
No SAV found	0	0	0	0	0	0	0	0	0		0%	